

Quantitative PCR in soil-transmitted helminth programs

Cools Piet, Verweij Jaco J, Ayana Mio, Albonico Marco, Ame Shaali, Dana Daniel, Keiser Jennifer, Mekonnen Zeleke, Sayasone Somphou, Vercruysse Jozef, Vlaminck Johnny and Levecke Bruno.

Joint meeting of the BSPP, ISP, BAVP and EVPC


4th of May 2018

Brussels, Belgium



Background

- Soil-transmitted helminths (STHs)
 - *Ascaris lumbricoides*
 - *Trichuris trichiura*
 - hookworm (*Necator americanus* & *Ancylostoma duodenale*)
- STHs > 3 million disability-adjusted life-years worldwide



Background

- Cornerstone control is mass drug administration (MDA)

population-level prevalence



time


Public health problem?
Which type of MDA?

Drugs working?
Program working?

Stop MDA?

Background

- Diagnostic tools essential in each decision step
- WHO: microscopic detection STHs eggs in stool Kato-Katz smear



Objectives




- Investigate applicability of quantitative PCR (qPCR) in STHs programs
 - Sensitivity
 - Drug efficacy
 - Classes of intensity of infection (low, moderate, high)
 - Pooling of stool

Methodology

- Drug efficacy trials (Tanzania, Laos, Ethiopia)
- School children baseline and follow-up visit
- Stool ethanol preservation, DNA extraction and qPCR for *A. lumbricoides*, *T. trichiura*, *N. americanus* and *A. duodenale*

Methodology

- Egg counts 4 microscopic methods
 - single Kato-Katz
 - Duplicate Kato-Katz
 - Mini-FLOTAC
 - FECPAK^{G2}



Methodology

- Sensitivity: composite reference method gold standard
- Classes intensity of infection: ROC analysis
- Drug efficacy:
 - Egg reduction rate (ERR) (KK) or genomic equivalents reduction rate (GRR) (qPCR)

$$= 100\% \times \left(1 - \frac{\text{arithmetic mean of group FEC at follow-up}}{\text{arithmetic mean of group FEC at baseline}} \right)$$

- WHO

	<i>A. lumbricoides</i>	<i>T. trichiura</i>	hookworm
Satisfactory	≥ 95	≥ 50	≥ 90
Doubtful	85 < ERR < 95	40 < ERR < 50	80 < ERR < 90
Reduced	≤ 85	≤ 40	≤ 80

- Individual samples vs pools of 10, 20 & 60

Findings: sensitivity


- qPCR is most sensitive method

	<i>Ascaris</i> n=540	<i>Trichuris</i> n=888	hookworm n=674
qPCR	90.0 [87.4; 92.4]	94.7 [93.2; 96.2]	92.0 [89.1; 93.9]
2x Kato-Katz	73.0 [69.3; 76.7]**	91.0 [89.0; 92.9]*	75.2 [72.0; 78.5]**
1x Kato-Katz	71.9 [68.0; 75.6]**	88.2 [86.0; 90.2]**	72.6 [69.1; 75.8]**
Mini-FLOTAC	63.3 [59.3; 67.4]**	91.6 [89.6; 93.4]*	73.9 [70.6; 77.2]**
FECPAK ^{G2}	58.9 [54.8; 63.1]**	60.0 [56.5; 63.1]**	52.4 [48.5; 56.2]**

* p<0.05 ** p<0.001


Findings: sensitivity

- qPCR is most sensitive method




Findings: classes of infection

Trichuris trichiura



Findings: classes of infection

Trichuris trichiura



Findings: drug efficacy

		Kato-Katz	qPCR
<i>Ascaris</i>	Ethiopia	99.9	99.1
	Laos	99.2	99.1
	Tanzania	96.7	96.8
<i>Trichuris</i>	Ethiopia	48.1	67.5
	Laos	40.5	-4.4
	Tanzania	-4.9	49.0
hookworm	Ethiopia	96.3	94.4
	Laos	96.2	94.6
	Tanzania	83.6	78.1

Findings: drug efficacy

		Kato-Katz	qPCR	
<i>Ascaris</i>	Ethiopia	99.9	99.1	normal
	Laos	99.2	99.1	
	Tanzania	96.7	96.8	
<i>Trichuris</i>	Ethiopia	48.1	67.5	doubtful
	Laos	40.5	-4.4	
	Tanzania	-4.9	49.0	
hookworm	Ethiopia	96.3	94.4	reduced
	Laos	96.2	94.6	
	Tanzania	83.6	78.1	

normal

doubtful

reduced




ASSESSING THE EFFICACY OF ANTHELMINTHIC
DRUGS AGAINST SCHISTOSOMIASIS
AND SOIL-TRANSMITTED HELMINTHIASES


Findings: pooling

- Pooling holds promise


Ascaris



Trichuris



hookworm



Mean # of genome copies from 10 individual samples

Conclusions

- Higher sensitivity
- Pooling promising and could reduce costs
- Monitor drug efficacy
- Cost-effectiveness analysis
- Place in STH program: low prevalence/intensity of infection